

2

OFFICE OF NAVAL RESEARCH

Grant N00014-90-J-1193

TECHNICAL REPORT No. 9

AD-A220 723

Intersubband Transitions in an Asymmetric Quantum Well with a Thin Barrier of a Delta-Function Potential

by

L. N. Pandey, Thomas F. George and M. L. Rustgi

Prepared for Publication

in

Journal of Applied Physics (Communication)

Departments of Chemistry and Physics  
State University of New York at Buffalo  
Buffalo, New York 14260

April 1990

DTIC  
ELECTE  
APR 23 1990  
S D  
Cc D

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

04 17 085

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/90/TR-9			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260			7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Grant N00014-90-J-1193		
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Intersubband Transitions in an Asymmetric Quantum Well with a Thin Barrier of a Delta-Function Potential					
12. PERSONAL AUTHOR(S) L. N. Pandey, Thomas F. George and M. L. Rustgi					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) April 1990	
				15. PAGE COUNT 15	
16. SUPPLEMENTARY NOTATION Prepared for publication in the Journal of Applied Physics (Communications)					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	QUANTUM WELL DELTA-FUNCTION POTENTIAL		
			INTERSUBBAND TRANSITIONS, OSCILLATOR STRENGTHS		
			THIN BARRIER; THEORY		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The changes in the bound-state energies and oscillator strength for intersubband transitions brought about by a thin barrier in the middle of an asymmetric quantum well are calculated, with a particularly close look at such changes as the middle barrier height approaches the bound-state energies. It is found that the oscillator strength goes through a slight change as the barrier height approaches the ground-state energy but an abrupt change when it approaches the excited bound-state energy. A suitable explanation for this change is provided. A similar tailoring of the intersubband transitions is also achieved by placing a delta-function potential in the vicinity of the middle of the well but without any abrupt change in the oscillator strength.					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

**Intersubband transitions in an asymmetric quantum well  
with a thin barrier or a delta-function potential**

L. N. Pandey, Thomas F. George and M. L. Rustgi  
Department of Physics & Astronomy and Chemistry  
Center for Electronic and Electro-optic Materials  
State University of New York at Buffalo  
Buffalo, New York 14260

The changes in the bound-state energies and oscillator strength for intersubband transitions brought about by a thin barrier in the middle of an asymmetric quantum well are calculated, with a particularly close look at such changes as the middle barrier height approaches the bound-state energies. It is found that the oscillator strength goes through a slight change as the barrier height approaches the ground-state energy but an abrupt change when it approaches the excited bound-state energy. A suitable explanation for this change is provided. A similar tailoring of the intersubband transitions is also achieved by placing a delta-function potential in the vicinity of the middle of the well but without any abrupt change in the oscillator strength.

1990 PACS Nos. : 73.20.Dx, 78.65.Fa, 73.60.Br



Accession For	
NTIS	CRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced <input type="checkbox"/>	
Justification _____	
By _____	
Distribution / _____	
Availability Codes	
Dist	Acquisition or Special
A-1	

In the last decade, enormous effort has been spent in understanding the physics of quantum well structures because of their applications to electronic and optoelectronic devices.<sup>1</sup> These structures can be fabricated by the molecular beam epitaxy technique, where the thicknesses and fractional molecular constituents of the layers can be controlled precisely and the interfaces sharply defined. The most widely used structure consists of alternate layers of  $Al_xGa_{1-x}As$  and  $GaAs$  where  $Al_xGa_{1-x}As$  has a direct band gap at the  $\Gamma$ -point for  $Al$  concentration  $x$  less than 0.4. The conduction and valence band differences in  $Al_xGa_{1-x}As$  and  $GaAs$  layers form the barriers and wells, respectively. A  $GaAs$  layer of appropriate thickness sandwiched between the two thick layers of  $Al_xGa_{1-x}As$  acts as a quantum well, which has different bound states depending upon the  $GaAs$  layer thickness and  $Al$  concentration  $x$  in the  $Al_xGa_{1-x}As$  layers. Recently, transitions between the bound states of the well have received considerable amount of theoretical and experimental attention because of the applicability to photoconductivity<sup>2</sup> and light absorption.<sup>3-6</sup> It has also been found that with suitably chosen parameters for the quantum well, the structure could be used for infrared detection.<sup>7-10</sup>

Very recently, Trzeciakowski and McCombe<sup>10</sup> performed a theoretical study to tailor the intersubband absorption by considering a thin layer of  $Al_yGa_{1-y}As$  in the middle of the well. They found that the presence of a thin barrier in the middle of the well shifts the ground-state energy of the well whereas the first-excited state remains practically unchanged. By changing the concentration  $y$  of  $Al$  in the middle layer, which determines the height of the barrier, they demonstrated that the energy separation between the ground state and the first-excited state could be changed without appreciably changing the oscil-

lator strength for the ground state to the first-excited state .

In the present Letter, we report the results of an investigation similar to that of Trzeciakowski and McCombe<sup>10</sup> but with a closer look at cases where the height of the middle barrier approaches the energy of the ground state and the first-excited state. A different pattern for the change in the oscillator strength as function of barrier height is found as the barrier height approaches the bound states. It is also found that a similar tailoring can also be achieved by changing the strength of a delta-function potential placed in the vicinity of the middle of the well without any appreciable change in the oscillator strength.

An asymmetric coupled well as shown in Fig. 1 has been considered for investigation. Some approximations such as neglecting the  $\Gamma$ -X mixing when Al concentration  $y$  is more than 0.4 and nonparabolicity effects<sup>11</sup> which affect the results slightly are made for simplification. We also take a constant effective mass ( $0.067m_0$ ,  $m_0$  being the rest mass of the electron) throughout the structure. The well extends from  $-d/2$  to  $d/2$  where  $d$  is taken to be 150 Å. The thin barrier in the middle extends from  $-b/2$  to  $b/2$  and three different values of  $b$  (5, 10 and 15 Å) are considered in our calculation. For the calculation with delta-function potential in the vicinity of the middle of the well, three different positions ( $c = 0, 5, 10$  Å) from the middle of the well are used. All the energies are measured from the bottom of the well. The energies of both the symmetric and antisymmetric bound states are given by the relation

$$\begin{aligned} & \sin^2 \frac{kd}{2} \left[ \frac{\bar{K}_L}{k} A_{11} + \frac{\bar{K}_L \bar{K}_R}{k^2} A_{12} + A_{21} + \frac{\bar{K}_R}{k} A_{22} \right] \\ & + \cos^2 \frac{kd}{2} \left[ -\frac{\bar{K}_L}{k} A_{11} + A_{12} + \frac{\bar{K}_L \bar{K}_R}{k^2} A_{21} - \frac{\bar{K}_R}{k} A_{22} \right] \end{aligned}$$

$$-\frac{1}{2} \sin kd \left[ \left( 1 - \frac{K_L K_R}{k^2} \right) A_{11} - \frac{K_L + K_R}{k} A_{12} - \frac{K_L - K_R}{k} A_{21} - \left( 1 - \frac{K_L K_R}{k^2} \right) A_{22} \right] = 0. \quad (1)$$

where  $k = \sqrt{\frac{2m^*E}{\hbar^2}}$ ,  $K_L = \sqrt{\frac{2m^*(V_L - E)}{\hbar^2}}$ ,  $K_R = \sqrt{\frac{2m^*(V_R - E)}{\hbar^2}}$ ,  $m^*$  is an effective mass, and  $A$  is a  $2 \times 2$  matrix which results from matching the wave functions and their derivatives at the interfaces of the middle barrier or at the delta-function potential.  $V_L$  and  $V_R$  are the depths of the well as shown in Fig. 1. For the middle barrier extending from  $-b/2$  to  $b/2$ ,  $A$  is given by

$$A_{11} = \cos kb \cos k_b b + \frac{1}{2} \left( \frac{k_b}{k} + \frac{k}{k_b} \right) \sin kb \sin k_b b. \quad (2a)$$

$$A_{12} = -\sin kb \cos k_b b + \left( \frac{k_b}{k} \cos^2 \frac{kb}{2} - \frac{k}{k_b} \sin^2 \frac{kb}{2} \right) \sin k_b b. \quad (2b)$$

$$A_{21} = \sin kb \cos k_b b + \left( \frac{k_b}{k} \sin^2 \frac{kb}{2} - \frac{k}{k_b} \cos^2 \frac{kb}{2} \right) \sin k_b b. \quad (2c)$$

$$A_{22} = \cos kb \cos k_b b + \frac{1}{2} \left( \frac{k_b}{k} + \frac{k}{k_b} \right) \sin kb \sin k_b b. \quad (2d)$$

when the barrier height  $V_b$  is less than  $E$ . The same expression can be used when  $V_b$  is greater than  $E$  by changing  $k_b$  to  $ik_b$ . Here  $k_b = \sqrt{\frac{2m^*(E - V_b)}{\hbar^2}}$ . For the delta-function potential located at a distance  $c$  from the middle of the well, the matrix  $A$  takes the form

$$A = \begin{pmatrix} 1 - \frac{\delta}{2k} \sin 2kc & -\frac{\delta}{k} \cos^2 kc \\ \frac{\delta}{k} \sin^2 kc & 1 + \frac{\delta}{2k} \sin 2kc \end{pmatrix}. \quad (3)$$

where  $\delta = \frac{2m^*\delta'}{\hbar^2}$ , with  $\delta'$  as the strength of the delta-function potential measured in  $\text{eV} \cdot \text{\AA}$ . It is clear from Eqs. (1)-(3) that for the case where  $V_b$  or  $\delta'$  is taken to be zero,  $A$  reduces to a unit matrix and Eq. (1) takes the form

$$\frac{K_L + K_R}{k} \cos kd - \left( 1 - \frac{K_L K_R}{k^2} \right) \sin kd = 0. \quad (4)$$

which is the condition for the bound states in an asymmetric well of width  $d$ . For the symmetric well, since  $K_L = K_R \equiv K$ , Eq. (4) reduces to

$$\cos kd - \frac{1}{2} \left( \frac{k}{K} + \frac{K}{k} \right) \sin kd = 0, \quad (5)$$

which is the condition for the bound states in a symmetric well of width  $d$ .

The oscillator strength for the dipole transition from a state  $m$  to a state  $n$  is given by

$$f_{mn} = \frac{2m_0(E_n - E_m)}{\hbar^2} | \langle n | x | m \rangle |^2, \quad (6)$$

where  $m_0$  is the rest mass of the electrons,  $E_n$  and  $E_m$  are the energies of the  $n$  and  $m$  states, respectively, and  $\langle n | x | m \rangle$  is the dipole matrix element.

Equations (1)-(3) can be used to find the symmetric and antisymmetric states of an asymmetric well, and the oscillator strength for the transition between these two states can be calculated by employing Eq. (6). Depending upon the width and depth of the well, if there is more than one excited state then a transition between the symmetric states is forbidden by selection rules. A transition from the ground state to the second-excited state is not allowed, whereas the transition from the first-excited state to the second-excited state is allowed if either of the states is populated by some means.

Figures (2) and (3) show our results for a well of width 150 Å with left and right walls 200 meV and 180 meV deep, respectively, and a thin barrier of width 5, 10 or 15 Å. Panel (a) of Fig. 2 shows the change in the ground-and the first-excited-state energies and the oscillator strength for the transition between them versus the barrier height in the vicinity of the ground state. The insert in the lower half of panel (a) displays the ground state and first-excited wave functions for the barrier height and width of 18 meV and 15

A, respectively. It is quite clear that the oscillator strength remains practically constant for the lower values of  $V_b$ . As  $V_b$  approaches the ground state, the change brought by the barrier in the ground-state wave function increases the oscillator strength slightly because the barrier remains practically unseen by the excited state, and also the wave function of the excited state vanishes in the vicinity of the middle of the well. The results in panel (b) of Fig. 2 display the changes due to the barrier height in the vicinity of the excited state. In this case the wave function of the ground state goes through a minimum in the barrier region, whereas the wave function of the excited state has an oscillation (due to the formation of an excited state in the small well formed by the excited state and the barrier) in that region, although the wave function still vanishes in the vicinity of the middle of the well (see the insert in the bottom half of panel (b) of Fig. 2). This change in the wave function results in a minimum for the oscillator strength, but the state energies are not affected. In Fig. 3, we show the results when  $V_b$  is above the energies of both states. A smooth variation of the oscillator strength as expected is found. The change in the ground-state energy and decrease of the oscillator strength results from the minimum of the ground-state wave function in the barrier region. A similar pattern is also observed for the transition from the first-excited state to the second-excited state but is not shown here in the interest of brevity.

Figure 4 shows the results for a delta-function potential of variable strength placed at 0, 5 or 10 Å away from the middle of the well. The oscillator strength in this case decreases smoothly, even for a small value of the strength of the delta-function potential, because the height of a delta-function potential extends to infinity.



In conclusion, we have carried out a theoretical investigation of the intersubband transitions in an asymmetric quantum well (1) by considering a thin barrier of variable height in the middle of the well or (2) by placing a delta-function potential of variable strength in the vicinity of the middle of the well. In the case of a thin barrier, significant changes in the oscillator strength are noticed as the barrier height approaches the bound state energies, whereas the energies of the states remained unaffected. It has also been found that the tailoring of intersubband transitions is also possible by changing the strength of a delta-function potential placed in the vicinity of the middle of the well without appreciably changing the oscillator strength.

One of the authors (MLR) is grateful to Dr. D.J. Nagel of the Naval Research Laboratory for providing the facilities to carry out a part of this work during the summer of 1989. This research was supported by the Office of Naval Research.

### References

1. L.L. Chang, L. Esaki and R. Tsu, Appl. Phys. Lett. **24**, 593 (1974).
2. B.F. Levine, K.K. Choi, C.G. Bethea, J. Walker and R.J. Malik, Appl. Phys. Lett. **50**, 1092 (1987).
3. M. Olszakier, E. Ehrenfreund, E. Cohen, J. Bajaj, G.J. Sullivan and D. Miller, Superlattices and Microstructures **5**, 283 (1989).
4. R. Cingolani, L. Tapfer, Y.H. Zhang, R. Muralidharan, K. Ploog and C. Tejedor, Phys. Rev. B **40**, 8319 (1989).
5. L.C. West and S.J. Singh, Appl. Phys. Lett. **46**, 1156 (1985).
6. G. Li, W. Wobbelare, D. Huang and H. Morkoc, Phys. Rev. B **39** (1989).
7. B.F. Levine, R.J. Malik, J. Walker, K.K. Choi, C.G. Bethea, D.A. Kleinman and J.M. Vandenberg, Appl. Phys. Lett. **50**, 273 (1987).
8. M.A. Kinch and A. Yariv, Appl. Phys. Lett. **55**, 2093 (1989).
9. R.L. Greene and P. Lane, Phys. Rev. B **34**, 8639 (1986).
10. W. Trzeciakowski and B.D. McCombe, Appl. Phys. Lett. **55**, 891 (1989).
11. D.F. Nelson, R.C. Miller and D.A. Kleinman, Phys. Rev. B **35**, 7770 (1987).

### Figure Captions

- Fig. 1. Schematic diagram of an asymmetric quantum well structure with a thin barrier or a delta-function potential. The parameters taken for the calculation are as follows:  $V_L = 200$  meV,  $V_R = 180$  meV,  $d = 150$  Å,  $b = 5, 10$  or  $15$  Å, and  $c$  (position of the delta-function potential away from the middle of the well) =  $0, 5$  or  $10$  Å.
- Fig. 2. Energies of the ground and first-excited states measured from the bottom of the well and their corresponding oscillator strength versus barrier height (i) in the vicinity of the ground state. [panel (a)] and (ii) in the vicinity of the first excited state [panel (b)]. The solid, dashed and dotted curves correspond to barrier widths of  $5, 10$  and  $15$  Å, respectively. The insert in the bottom half of panel (a) displays the ground-state (solid line) and first-excited-state (dotted line) wave functions from  $-100$  to  $100$  Å spatial dimension for a  $18$  meV barrier height and  $15$  Å barrier width. In panel (b) the wave functions are shown for a  $64$  meV barrier height and  $15$  Å barrier width.
- Fig. 3. Ground and first-excited-state energies and oscillator strength as in Fig. 2 but for a barrier height above the first-excited state.
- Fig. 4. Energies of the ground and first-excited states measured from the bottom of the well and their corresponding oscillator strengths as a function of the strength of the delta-function potential. The solid,

dashed and dotted curves correspond to  $c = 0.5$  and  $10 \text{ \AA}$ , respectively.

Exactly the same results are found for  $c = -5$  and  $-10 \text{ \AA}$ .

Fig. 1




Fig. 2





Fig. 3





Fig. 4





TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. J. H. Weaver  
Department of Chemical Engineering  
and Materials Science  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. A. Reisman  
Microelectronics Center of North Carolina  
Research Triangle Park, North Carolina  
27709

Dr. M. Grunze  
Laboratory for Surface Science and  
Technology  
University of Maine  
Orono, Maine 04469

Dr. J. Butler  
Naval Research Laboratory  
Code 6115  
Washington D.C. 20375-5000

Dr. L. Interante  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Irvin Heard  
Chemistry and Physics Department  
Lincoln University  
Lincoln University, Pennsylvania 19352

Dr. K.J. Klaubunde  
Department of Chemistry  
Kansas State University  
Manhattan, Kansas 66506

Dr. C. B. Harris  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. F. Kutzler  
Department of Chemistry  
Box 5055  
Tennessee Technological University  
Cookeville, Tennessee 38501

Dr. D. DiLella  
Chemistry Department  
George Washington University  
Washington D.C. 20052

Dr. R. Reeves  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Steven M. George  
Stanford University  
Department of Chemistry  
Stanford, CA 94305

Dr. Mark Johnson  
Yale University  
Department of Chemistry  
New Haven, CT 06511-8118

Dr. W. Knauer  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. Murday  
Naval Research Laboratory  
Code 6170  
Washington, D.C. 20375-5000

Dr. J. B. Hudson  
Materials Division  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Theodore E. Madey  
Surface Chemistry Section  
Department of Commerce  
National Bureau of Standards  
Washington, D.C. 20234

Dr. J. E. Demuth  
IBM Corporation  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. M. G. Lagally  
Department of Metallurgical  
and Mining Engineering  
University of Wisconsin  
Madison, Wisconsin 53706

Dr. R. P. Van Duyne  
Chemistry Department  
Northwestern University  
Evanston, Illinois 60637

Dr. J. M. White  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. D. E. Harrison  
Department of Physics  
Naval Postgraduate School  
Monterey, California 93940

Dr. R. L. Park  
Director, Center of Materials  
Research  
University of Maryland  
College Park, Maryland 20742

Dr. W. T. Peria  
Electrical Engineering Department  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson  
Department of Metallurgy and  
Materials Science  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. S. Sibener  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Arnold Green  
Quantum Surface Dynamics Branch  
Code 3817  
Naval Weapons Center  
China Lake, California 93555

Dr. A. Wold  
Department of Chemistry  
Brown University  
Providence, Rhode Island 02912

Dr. S. L. Bernasek  
Department of Chemistry  
Princeton University  
Princeton, New Jersey 08544

Dr. W. Kohn  
Department of Physics  
University of California, San Diego  
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Richard Colton  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Dan Pierce  
National Bureau of Standards  
Optical Physics Division  
Washington, D.C. 20234

Dr. R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, California 90024

Dr. R. P. Messmer  
Materials Characterization Lab.  
General Electric Company  
Schenectady, New York 22217

Dr. Robert Gomer  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Ronald Lee  
R301  
Naval Surface Weapons Center  
White Oak  
Silver Spring, Maryland 20910

Dr. Paul Schoen  
Code 6190  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. John T. Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene  
Code 5230  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. L. Kesmodel  
Department of Physics  
Indiana University  
Bloomington, Indiana 47403

Dr. K. C. Janda  
University of Pittsburgh  
Chemistry Building  
Pittsburg, PA 15260

Dr. E. A. Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, North Carolina 27514

Dr. Adam Heller  
Bell Laboratories  
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann  
Department of Chemistry  
University of Southampton  
Southampton SO9 5NH  
UNITED KINGDOM

Dr. H. Tachikawa  
Chemistry Department  
Jackson State University  
Jackson, Mississippi 39217

Dr. John W. Wilkins  
Cornell University  
Laboratory of Atomic and  
Solid State Physics  
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis  
Department of Physics  
University of California  
Irvine, California 92664

Dr. D. Ramaker  
Chemistry Department  
George Washington University  
Washington, D.C. 20052

Dr. J. C. Hemminger  
Chemistry Department  
University of California  
Irvine, California 92717

Dr. T. F. George  
Chemistry Department  
University of Rochester  
Rochester, New York 14627

Dr. G. Rubloff  
IBM  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. Horia Metiu  
Chemistry Department  
University of California  
Santa Barbara, California 93106

Dr. W. Goddard  
Department of Chemistry and Chemical  
Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. P. Hansma  
Department of Physics  
University of California  
Santa Barbara, California 93106

Dr. J. Baldeschwieler  
Department of Chemistry and  
Chemical Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. J. T. Keiser  
Department of Chemistry  
University of Richmond  
Richmond, Virginia 23173

Dr. R. W. Plummer  
Department of Physics  
University of Pennsylvania  
Philadelphia, Pennsylvania 19104

Dr. E. Yeager  
Department of Chemistry  
Case Western Reserve University  
Cleveland, Ohio 41106

Dr. N. Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, Pennsylvania 16802

Dr. Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. A. Steckl  
Department of Electrical and  
Systems Engineering  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. G.H. Morrison  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853